

1.5×9 mm, 1.5×20 mm, 2.5×25 mm, 2.75×15 mm, 2.75×18 mm, and 2.75×23 mm non-compliant (NC) balloons, followed by a 2.5×20 mm scoring balloon and a 2.5×15 mm cutting balloon. Although the cutting balloon ruptured at 14 atm, adequate plaque modification could not be achieved. Given the development of a limited dissection with preserved TIMI grade 3 flow and the absence of symptoms, the procedure was terminated, and rotational atherectomy was planned.

Second Session (April 8, 2026): The lesion was successfully crossed with the support of a microcatheter. Following predilation, rotational atherectomy was performed using a 1.75-mm burr. Subsequent high-pressure dilations with a 3.25-mm cutting balloon, a 3.0×12 mm NC balloon inflated up to 34 atm, and a 3.5×15 mm NC balloon inflated up to 26 atm failed to achieve adequate lumen expansion. The lesion was considered balloon-undilatable, and repeat intervention with either intravascular lithotripsy (IVL) or an OPN balloon was planned. Following the procedure, new-onset atrial fibrillation developed, and oral anticoagulation therapy was initiated.

Third Session (April 27, 2026): The previously rotatable resistant lesion was re-crossed with a guidewire. A 2.5×15 mm OPN balloon was inflated up to 40 atm, followed by additional high-pressure dilation with a 3.0×20 mm NC balloon. Despite these interventions, persistent balloon waist was observed, and adequate lumen expansion could not be achieved. No perforation, no-reflow phenomenon, or hemodynamic instability occurred during the procedure. Given the failure to achieve technical success despite sequential plaque modification techniques, further attempts at revascularization were abandoned, and optimal medical therapy was pursued. The patient was discharged without cardiac symptoms.

Conclusion: In severely calcified CTO lesions, technical success may not always be achieved despite the sequential use of advanced plaque modification techniques, including rotational atherectomy, cutting balloons, and ultra-high-pressure balloons. This case highlights that balloon-undilatable calcified CTO lesions represent one of the most challenging scenarios in interventional cardiology. Intravascular imaging-guided procedural planning and the timely use of emerging calcium-modification technologies, such as IVL, may represent potential strategies to improve procedural success in similar cases.

SS-17

Rotablation-Assisted Complex Left Main Coronary Trifurcation Intervention in Severe Calcification: A Successful IVUS-Guided Complex Percutaneous Coronary Intervention

Yasin Aktaş, Gülşah Karaduman, Hakan Erkan

Bursa State Hospital, Bursa, Türkiye

Introduction and Aim: Left main coronary artery (LMCA) trifurcation lesions, particularly in the presence of severe calcification, represent some of the most complex interventions in interventional cardiology. In patients who are unsuitable for surgical revascularization or decline surgery, rotational atherectomy-assisted percutaneous coronary intervention may provide a successful alternative.

Case Report: A 69-year-old man was admitted with non-ST-segment elevation myocardial infarction and acute decompensated heart failure. Coronary angiography revealed a severely calcified Medina 0,1,1 LMCA trifurcation lesion, accompanied by a chronic total occlusion of the distal right coronary artery. Although surgical revascularization was recommended by the Heart Team, the patient declined surgery, and a high-risk percutaneous intervention was planned. For lesion preparation, rotational atherectomy was performed using a 1.5-mm burr, followed by implantation of a drug-eluting stent from the LMCA into the circumflex artery. During the procedure, impairment of LAD flow developed and was successfully managed using the T and small protrusion (TAP) technique, followed by final kissing balloon inflation and proximal opti-

mization technique. Intravascular ultrasound confirmed adequate stent expansion and complete apposition. The procedure was completed without complications, with restoration of TIMI grade 3 flow on final angiography. Follow-up angiography demonstrated patent stents, while the patient showed a marked improvement in functional capacity and an increase in left ventricular ejection fraction during follow-up.

Conclusion: This case demonstrates that, in patients with severely calcified LMCA trifurcation lesions, a strategy combining rotational atherectomy, an appropriate bifurcation technique, and IVUS-guided stent optimization may provide a safe and successful treatment option for high-risk patients.

Keywords: Left main coronary artery trifurcation, rotational atherectomy, intravascular ultrasound (IVUS).

SS-18

Management of Aorto-Ostial Stent Protrusion Using the Side Flap Technique in Acute Coronary Syndrome Developing During Chemotherapy: A Case Report

Yusuf Can, Ömer Faruk Erkan, Sabri Çiftçi

Department of Cardiology, Sakarya University Faculty of Medicine, Sakarya, Türkiye

Introduction and Aim: Aorto-ostial stent implantation is a technically challenging percutaneous coronary intervention due to anatomical considerations and difficulties in achieving optimal stent positioning. Stent protrusion into the aorta may make subsequent re-entry into the stent lumen difficult and compromise adequate catheter support during further interventions. The Side Flap technique is an alternative approach that enables the creation of a new functional lumen by passing through the side wall of the stent in selected cases where re-entry into the original stent lumen is not feasible. In this case, we present the successful management of aorto-ostial stent protrusion using the Side Flap technique during coronary intervention performed for chest pain and elevated troponin levels that developed during chemotherapy.

Case Report: A 65-year-old man with a history of hypertension and coronary artery disease, including previous coronary stent implantation, was admitted for evaluation of chest pain and elevated troponin levels that developed during chemotherapy for lung cancer. Electrocardiography showed sinus rhythm. Echocardiography revealed a left ventricular ejection fraction of 60%, mild mitral and tricuspid regurgitation, and grade I diastolic dysfunction. Coronary angiography demonstrated an 80% stenosis at the LAD ostium, a patent stent in the mid-LAD segment, a 30% stenosis in the CX-OM2 segment, and severe stenoses in the ostial (90%) and mid (95%) segments of the nondominant RCA. Percutaneous coronary intervention (PCI) to the LAD was planned, and the left main coronary artery (LMCA) was engaged with a 7-French 3.5 guiding catheter. After wiring the LAD and CX, balloon angioplasty of the LAD was performed using a 2.0×25 mm balloon. Subsequently, a 3.5×38 mm drug-eluting stent (DES) was deployed from the LMCA ostium into the LAD. The stent was assessed in different angiographic projections and positioned with 1–2 struts extending into the aorta. During stent deployment, the patient took a deep inspiration, resulting in approximately 10 mm of stent protrusion into the aorta. Because of the stent protrusion into the LMCA ostium, inability to re-cross the stent lumen with a guidewire, and compromised CX flow, the guidewire was advanced from the LMCA into the LAD by passing through the edge of the protruding stent struts. At this stage, the Side Flap technique was performed using a 3.5×10 mm balloon, resulting in lateral displacement of the protruding stent segment and creation of a functional lumen. Proximal optimization technique (POT) was then performed in the LMCA using a 5.0×8 mm balloon. A 3.5×23 mm DES was implanted between the two stents in the LAD. The CX was rewired with a floppy guidewire, and the stent struts were dilated using a 2.5×15 mm balloon. A 3.0×34 mm DES was subsequently implanted in the CX using the TAP technique. After kiss-